

HIV/AIDS

HIV/AIDS Precautions - Workplace Concerns

On this page

[What are some basic facts about HIV?](#)

[Who are the workers at risk?](#)

[How is HIV transmitted?](#)

[How is transmission in the workplace prevented?](#)

What are some basic facts about HIV?

The Human Immunodeficiency Virus (HIV) is a virus that infects the immune system. Acquired Immune Deficiency Syndrome (AIDS) is the most advanced stage of the HIV infection. HIV causes the immune system to become vulnerable to other infections. There is presently no cure or vaccine for HIV. It can take, on average, up to 10 years for the initial HIV infection to progress to AIDS without treatment.

Please also see the OSH Answers document [HIV/AIDS in the Workplace](#) for more information.

How is HIV transmitted?

In the workplace, the most likely method HIV is transmitted is an injury from a contaminated needle or other sharp objects. Other ways the human immunodeficiency virus (HIV) can pass from one person to another in the following ways:

- by unprotected sexual intercourse with infected persons
- via transfusion of infected blood or blood products
- from an infected mother to her infant before or during birth, or through breastfeeding
- via organ transplant from an infected donor

HIV is not found in saliva, vomit, feces, nasal secretions, tears or urine unless these fluids are visibly contaminated with blood.

HIV is not spread by casual contact. There is no risk of becoming infected with HIV by working on the same assembly line, using the same equipment, sharing locker rooms or toilet facilities or being in the same office as someone with HIV infection or AIDS.

Who are the workers at risk?

All workers who are in contact with contaminated blood or other specific body fluids are at risk. Exposure to HIV in the workplace occurs through:

- skin and mucous membrane contact with blood and other body fluids of an infected person
 - incidents with needles or other sharp instruments contaminated with the blood of an infected person
-

How is transmission in the workplace prevented?

The Centers for Disease Control recommend using [routine practices](#) to protect workers at risk from HIV exposure. This approach stresses that all situations involving contact with blood and certain other body fluids present a risk. Routine practices outline the use of barriers to prevent workplace exposure to HIV and other viruses. These barriers include the use of:

- engineering controls such as retractable needles
 - safe work practices and administrative controls
 - protective equipment such as gloves, gowns or aprons, masks, and protective eye wear.
-

Fact sheet confirmed current: 2017-03-01

Fact sheet last revised: 2022-12-23

Disclaimer

Although every effort is made to ensure the accuracy, currency and completeness of the information, CCOHS does not guarantee, warrant, represent or undertake that the information provided is correct, accurate or current. CCOHS is not liable for any loss, claim, or demand arising directly or indirectly from any use or reliance upon the information.